

# Parallel versus serial processing: new vistas on the distributed organization of the visual system

Jean Bullier and Lionel G Nowak

Cerveau et Vision, INSERM 371, Bron/Lyon, France

Recent functional studies question the validity of the hierarchical model of organization for processing visual information in cortical areas. The results of these studies suggest that beyond the primary visual cortex (V1), information is not serially processed through successive cortical areas, but that it is simultaneously processed in several areas. The idea that visual information is functionally segregated into different, parallel channels as it circulates through V1 and V2 towards V4 and the middle temporal visual area is also challenged by recent studies that report a smaller degree of functional specialization within the visual areas than previously thought.

Current Opinion in Neurobiology 1995, 5:497–503

## Introduction

Since the early days of single-unit recording in visual cortex, the models of cortical processing have vacillated between favoring a serial (hierarchical) or a parallel organization. The hierarchical model, originally proposed by Hubel and Wiesel, was shaped by the current opinion of the time, which postulated that several successive processing steps should separate sensation from perception (see [1\*]). In the cat, these successive steps were represented by cortical areas 17, 18 and 19, and by the simple, complex and hypercomplex functional classes of neurons. Thus, the serial model has two main characteristics: first, processing is accomplished through a cascade of effectors (neuron types or cortical areas), and second, progression in the hierarchy corresponds to an increasing 'complexity' of the effectors.

With the discovery in the 1970s that distinct functional types of neurons exist in the retina and the lateral geniculate nucleus (LGN) of the cat, the parallel model of organization gained recognition. X-, Y- and W-type of retinal ganglion cells appeared to provide the major inputs to different cortical areas [2], and lesions or inactivation (e.g. by cooling) of area 17 did not silence neurons in area 18, as would be expected if these areas operated in an hierarchical cascade [2,3]. Parallel models placed the emphasis on independent and simultaneous processing by modules that are specialized for different aspects of the visual stimulus.

The shift of interest from cat to monkey cortical anatomy and physiology in the 1980s brought back the concept of a hierarchical organization of cortical areas. This followed the early observation of Rockland

and Pandya [4] that different types of cortico-cortical connections, later called feedforward and feedback, can be distinguished by the laminar distributions of the parent neurons and target terminal arborizations. Identification of feedforward and feedback connections led to a hierarchical classification of cortical areas by the following criterion: higher-order areas receive feedforward connections and send feedback connections to lower-order areas [5].

At the same period, the idea that magnocellular and parvocellular layers of the LGN process different types of information in parallel became prevalent. In cortex, cytochrome oxidase staining revealed the presence of modules — blobs in cortical area V1 and bands in cortical area V2 — that appeared to operate in parallel.


All these findings lead us to a textbook picture of the organization of the visual system (Fig. 1), which is a hybrid of serial and parallel processing: namely, parallel channels operating within a hierarchy of cortical areas. This review examines recent findings concerning parallel and hierarchical processing in visual cortex, and it will become apparent that this hybrid model does not correspond with reality.

## Is there a hierarchy of visual cortical areas?

The major weakness of the hybrid hierarchical/parallel model is that it is based solely on anatomical observations. Functional studies provide little support for a hierarchical (serial) organization of extrastriate cortical areas.

### Abbreviations

K—koniocellular; LGN—lateral geniculate nucleus; M—magnocellular; MT—middle temporal; P—parvocellular; PMLS—postero-medial lateral suprasylvian; V1—primary visual cortex.



**Fig. 1.** Cartoon representation of the hybrid hierarchical/parallel model. For a more exact and detailed description, see [5]. Neurons of the parvocellular (P) and magnocellular (M) layers of the LGN give rise to parallel channels that circulate through area V1 and extrastriate cortex towards higher-order areas. The P pathway is supposed to give rise to two channels, one for the processing of color and the other devoted to form. These channels are associated to different cortical territories identified by cytochrome-oxidase (C-O) histochemistry and lead to V4 and intertemporal (IT) cortex. The M channel is supposed to circulate in parallel to the P channels, to be devoted to motion processing, and to innervate area MT, the parietal and upper superior temporal sulcus (STS) cortex. 4B, layer 4B of V1.


### Serial processing: effects of lesions or inactivation of lower-order areas

A serial organization assumes that information is processed at each step before being sent to the next level up. One way to test this mode of operation is to lesion or inactivate a lower-order area to determine whether this silences neurons in higher-order areas. This has only been tested in the V1 area of monkeys, which, when inactivated, silences areas V2, V3, V4 and the inferotemporal cortex [3,6–10]. These electrophysiological results are complemented by behavioural data: lesioning area V1 produces a marked visual deficit (see [11]). Interestingly, it has been shown that with re-training, these V1-lesioned animals can recover some form of vision [12], even though they appear to have no 'conscious' perception of the stimulus they can detect [13\*].

A possible explanation for this form of residual vision is that there appears to be some remaining neural activity in other extrastriate cortical areas. Lesioning or inactivation of area V1 decreases the response and lowers the selectivity of neurons in area V3A, the middle temporal (MT) area and the superior temporal polysensory (STP)

area, but does not silence them [14,15]. A similar finding has been reported in humans who have damaged V1 areas; despite the fact that no activity is recorded in the V1 area of these patients, a cortical region corresponding to area MT in monkeys appears to be active [16]. This residual activity in humans may be the functional basis of the phenomenon of blindsight [17,18].

Lesion and inactivation studies have thus confirmed the special status of V1 as the first relay for visual information coming from the thalamus and directed toward the inferotemporal cortex (see Fig. 1). However, this is not the case for the transfer of information directed toward the parietal cortex (see Fig. 1). Beyond V1, area V2 is the second relay of visual information in the hierarchy. Does this mean that inactivation of V2, like that of V1, will silence other higher-order extrastriate areas? This question has not been addressed directly. However, lesioning V2 in monkeys had little consequence on the visual abilities of the animal and only produced a very specific deficit in the detection of pop-out stimuli based on orientation differences [11]. The specificity of the deficit caused by lesioning V2 contrasts with the almost total blindness caused by lesioning V1 [11,13\*].



**Fig. 2.** Latency distribution of neurons in different layers of V1 and V2 responding to visual stimulation. The grey horizontal rectangles represent the range of latencies corresponding to the 25th–75th centile of activated neurons (vertical bars correspond to the median). The small horizontal bars extruding from the left and right side of the rectangles correspond to the latencies for the 10th and 90th centile of activated neurons, respectively. Note that the order of activation does not follow a serial V1-to-V2 order, but is heavily influenced by the type of LGN input (magnocellular to 4C $\alpha$  and parvocellular to 4C $\beta$ ). See [30\*,31\*] for further details.

This suggests that complete inactivation of V2 will not silence neurons in other higher-order cortical areas, and that, based on the criterion mentioned above, area V2 is not a lower-order area than V3, MT or V4, as proposed by the hierarchical model.

The role of V1 as the major relay of visual information is specific to primates because in non-primate species, lesioning or inactivation of area V1 does not silence other cortical areas. For example, in cats, although inactivation of V1 strongly decreases the responses of most neurons in area 21a, other neurons are little affected [19]. It also has been known for some time that lesioning cat V1 does not silence the responses of neurons of the suprasylvian sulcus [20,21] and that inactivation of V1 does not silence neurons in area 18 [22,23].

The mild effects of area V1 lesions on the activity in other cortical areas of the cat are in contradiction with the predictions of the hierarchical model because, in cats, as in monkeys, area VI occupies the lowest level in the hierarchy [5]. These electrophysiological results are paralleled by the small and specific visual deficits observed in cats after such lesions [24,25]. As has been argued elsewhere [3], the devastating effects of V1 lesions on vision observed in monkeys and humans is a feature unique to primates, which may be related to the more focussed LGN projections to the cortex.

To summarize, lesion or inactivation studies confirm the serial or hierarchical model of cortical processing only in the case of V1 in primates. In other cortical areas and in other orders, it appears that the hierarchical ranking of cortical areas based on the anatomy does not constitute a proper model of the functional visual system.

#### Serial processing: timing

Another way to test whether cortical areas process information in a serial fashion is to measure the latencies with which neurons in different cortical areas respond to visual stimulation. If information is indeed processed serially, then the latencies of higher-order areas should

be longer than those of the lower-order areas that drive them. Latencies are measured by flashing small spots of light at or by moving stimuli across the receptive fields of visual neurons. Such experiments revealed that there is a very large range of latencies within each cortical area, and that the timing of the latencies overlaps extensively across different areas [26–29].

The latencies in both area V1 and area V2 of the macaque monkey have recently been measured under the same stimulus conditions [30\*,31\*]. The results show a large overlap of the latency distributions, with V1 neurons activated, on average, 10 ms earlier than neurons in V2. When the neuronal populations of the various channels that stream through V1 and V2 are analyzed separately (see below), it is found that the serial order between V1 and V2 is not the major determinant of the latency differences observed between these areas. Figure 2 illustrates the distribution of latencies in different layers of V1 and different cytochrome oxidase bands in V2, arranged in increasing order (see [30\*,31\*]). This demonstrates that the earliest activated neuronal populations are in layers 4C $\alpha$  and 4B of V1 and the thick cytochrome oxidase bands of V2, all territories that are almost exclusively under the control of the magnocellular layers of the LGN (see Fig. 1). After most of the magnocellular-driven activity has left V2 through projection neurons in the thick cytochrome oxidase bands, layer 4C $\beta$  is activated by LGN neurons of the parvocellular layers. This is followed by the bulk of responses in the supragranular layers of V1 and the thin cytochrome oxidase bands of V2.

These results demonstrate that latencies in V1 and V2 overlap extensively and that the latency difference between magnocellular and parvocellular-driven populations is larger (20 ms) than the latency difference between V1 and V2 neurons (10 ms). Although neurons in V2 depend upon V1 for their activation [6,7], there are many neurons in V2 that are active earlier than some V1 neurons; therefore, these 'early' V2 neurons are in a position to drive some of the 'later' V1 neurons. The work of Sandell and Schiller [32] on the inactivation of

V2 confirms that some V1 neurons depend upon V2 for their visual responses.

A few studies have been devoted to the latencies of neurons in area MT. Raiguel *et al.* [26] found that the median latency of MT neurons is 9 ms longer than that of V1 neurons, and that the shortest latencies in area MT are even shorter than those in V1 or V2. Measuring the latency of the average post-stimulus time histogram of neurons in areas MT and V1, Maunsell [33] concluded that activation of area MT is delayed by 11 ms with respect to activation of the transient neurons in V1. It is interesting to compare these results with those measuring corresponding populations in V1 and V2: the latency difference between the shortest (10th centile) latencies in layer 4C $\alpha$  in V1 and the thick cytochrome oxidase bands of V2 (10th centile) is 16 ms (Fig. 2). This suggests that MT neurons are activated at the same time as, and possibly earlier than, the neurons in the thick cytochrome oxidase bands of V2, which is in keeping with the results of Raiguel *et al.* [26] described above. Thus, neurons of the 'second' (V2) and 'fifth' (MT) stages of the proposed hierarchical arrangement of extrastriate areas are activated simultaneously. Therefore, it seems unlikely that MT neurons depend on input from V2 in the same way that neurons in V2 depend upon input from V1. In other words, there is no physiological evidence that MT is a higher-order area than V2.

Measuring latencies in cortical areas of the cat visual system lead to a similar conclusion [34\*]. Despite its position as a second-order area in the cortical hierarchy, area 18 contains the earliest activated cortical neurons, some 5 ms earlier than neurons of area 17 [34\*]. This probably results from the fact that area 18 is mostly driven by Y cells, which have shorter latencies than X cells in the retina [35]. Thus, in the cat visual cortex, as in the monkey visual cortex, the type of thalamic afferent (X/Y or magnocellular/parvocellular) appears to be a stronger determinant of the latency with which a cortical neuron responds to activation than the supposed rank of the cortical area it belongs to.

In addition, the latencies of activation in the cat visual cortex clearly differ from those predicted by the hierarchical ranking of areas. In the hierarchical scheme, the order of activation should be areas 17, 18, 19, PMLS; however, the order given by the latencies is 18, 17, PMLS, 19 [34\*]. This means that the hierarchical scheme is of little help for predicting the temporal sequence of activation of cortical areas.

### Is processing done in parallel?

There is strong anatomical and physiological evidence that three separate channels transfer information through the LGN toward visual cortex. These are called M (for magnocellular), P (for parvocellular) and K (for koniocellular) channels. It has been demonstrated by electrophysiological and lesion studies that, although

most stimuli activate all three channels, information about color and high spatial frequency is carried by the P channel, and that information about high temporal frequency and low contrast are carried by the M channel [36–40].

It has previously been suggested that the P and M subcortical channels remain separate in cortex [5,41]. The P pathway was supposed to separate into two branches through areas V1 and V2 towards area V4. The first branch, circulating through layer 4C $\beta$ , the cytochrome oxidase blobs in V1 and the thin cytochrome oxidase bands in V2, was supposed to carry color information. The other branch, passing through layer 4C $\beta$ , the interblob regions in V1 and the pale cytochrome oxidase bands in V2, was associated with the processing of form. The M pathway was thought to circulate almost exclusively through layers 4C $\alpha$  and 4B in V1 and the thick cytochrome oxidase bands in V2, to activate area MT and to be devoted to the processing of motion information. The concept of functional parallel pathways in areas V1 and V2 (Fig. 1) is, therefore, based on two assumptions: first, that pathways remain separate in areas V1 and V2, and second, that the channels in V1 and V2 and extrastriate areas beyond V1 and V2 are specialized processors for certain aspects of the visual stimuli.

### Are M and P pathways segregated in areas V1 and V2?

Recent anatomical studies have demonstrated that there is far more convergence of the M and P pathways in V1 and V2 than previously thought. M, P and K channels converge in the cytochrome oxidase blobs, and the M and P channels converge in the interblobs of V1 [42,43,44\*\*]. In V2, extensive interconnections between the different cytochrome oxidase bands have been demonstrated [45\*\*,46\*\*].

Physiological studies using selective inactivations of M and P layers in the LGN confirm that many neurons of V1 receive converging information from M and P pathways [47,48\*]. The M and P inputs to the interblobs in V1 may explain why many neurons in pale cytochrome oxidase bands in V2 respond to colored stimuli [49,50\*\*] and at low luminance contrast [50\*\*], and why neurons in area V4 appear to be equally driven by M and P neurons of the LGN [51\*\*]. The similar responses of neurons in V4 and MT neurons to fast moving stimuli [52\*] may also result from the strong M channel input into V4.

On the other hand, despite the evidence of a P-dominated layer 4A input to the thick cytochrome oxidase bands and strong interconnections of the bands in V2 [45\*\*], the concept of a relatively pure M channel circulating through layers 4C $\alpha$  and 4B in V1 and the thick cytochrome oxidase bands of V2 toward MT has been confirmed by physiological studies [52\*,53–55]. Responses of MT neurons are little affected by the inactivation of P layers [53]. MT neurons give good

responses to flicker and fast moving stimuli and show high contrast sensitivity [52•] and poor responses to chromatic contrast [54,55], all properties expected from neurons of the M channel. The results of latency studies described above (see also Fig. 2) provide evidence that, in addition, the M pathway is characterized by early activation and fast information transfer.

#### Functional specialization of channels and cortical areas

One of the strong arguments in favor of a specialized color channel through the cytochrome oxidase blobs in V1 and the thin bands in V2 was the presence of numerous color-selective neurons among the non-orientation selective cells in the blobs in V1 and the thin bands in V2 [41,56,57]. It is clear, however, that color-selective neurons are also present in large numbers in the interblobs in V1 [56,58] and in the pale bands in V2 [50••].

When the receptive field properties of neurons in the different cytochrome oxidase bands in V2 are measured quantitatively, it is found that all types of selectivities are found with varying proportions in each set of cytochrome oxidase bands. Thus color-selective neurons are more numerous in thin and pale cytochrome oxidase bands [49,50••], and orientation-selective neurons are found in larger numbers in thick and pale bands [50••,59]. Direction-selective neurons are more numerous in thick cytochrome oxidase bands ([49,50••]; see, however, [59]). Thick bands also contain larger proportions of depth-selective neurons, as originally proposed by Hubel and Livingstone [41]. However, the earlier report that end-stopped cells are concentrated in the pale bands [41] has not been confirmed [50••,59].

Thus, recent reports do not confirm the original view that form, color, motion and depth are processed in parallel channels in areas V1 and V2. Beyond V2, areas V4 and MT are usually presented as examples of functional specialization for motion and color, as evidenced by the presence of numerous direction-selective neurons in MT and color-selective neurons in V4. A recent report confirms that neurons in MT show little selectivity for color [54]. However, as mentioned above, the responses of neurons in V4 and MT to moving stimuli appear similar in terms of velocity selectivity [52•].

The results of lesion studies suggest that areas V4 and MT are not specialized processors for color, form and motion. Long-term effects of lesions in area MT and the surrounding cortex in the superior temporal sulcus on motion perception are rather mild [60,61•]. In a similar way, lesioning area V4 does not lead to a long-term impairment of color and shape discrimination [60,62]. This demonstrates that motion, color and shape information can be processed in the absence of areas V4 and MT, and that these areas can no longer be regarded as the sole specialized processors for these

attributes of visual stimuli. It should be noted, however, that the short-term effects of lesions on behavior are much stronger [60,63]. The significance of such rapid post-lesion recovery remains to be determined. To circumvent these difficulties related to post-lesion recovery, it is possible to use reversible inactivation methods to test the participation of a given cortical area to visual behaviour. Using such methods, Lomber, Payne and co-workers [64••] have demonstrated a total but reversible deficit in the identification of visual cues masked by moving visual noise when the lateral suprasylvian (LS) area of the cat cortex was reversibly inactivated.

#### Conclusions

Recent work on the visual system suggests that the organization of cortical areas is neither serial nor parallel. Beyond the well known fact derived from lesion studies that area V1 is a bottleneck for visual information in primates, there is no strong evidence from functional studies that visual information is processed serially through a hierarchy of extrastriate cortical areas. Instead, measurements of visual latencies in different cortical areas suggest two characteristics of processing visual information: first, that the order of activation does not necessarily follow the order suggested by the anatomically defined hierarchy of cortical areas; and second, that there is much more *simultaneity* of processing in the different visual areas than would be expected from a strictly serial organization. This simultaneity suggests that not only feedforward but also feedback connections may play a role in establishing some receptive field properties and, ultimately, in perception, as suggested by recent theoretical studies [65–67]. To further characterize the nature of the information transfers through these connections, one could inactivate specific extrastriate visual areas and measure the responses of neurons in the connected areas. Such studies would help us understand the complexity of the flow of information in visual cortex.

The concept of parallel organization, which implies a segregation of channels and a specialization of cortical areas, has also been challenged by recent studies. First, channels kept separate at the level of the retina and thalamus appear to *converge* on the same cells in area VI and beyond, in the occipito-temporal pathway. Second, information about color, motion or form appears to be distributed among different cortical areas. This *distributed* processing is confirmed by functional studies in humans, which show that not only one, but several cortical regions are specifically activated by one of the parameters (i.e. color, or motion, or form) of the stimulus [68,69]. It is currently postulated that it is through the interactions between these modules that perceptual decisions can be made [65–67,70].

## Acknowledgements

We thank Naura Chounlamountri for help in preparing this manuscript. We acknowledge support from the following sources: Ministère de l'Éducation Nationale et de la Culture (Action Thématique Sciences Cognitives), from the Programme Cognisciences of the CNRS (Perception et Reconnaissance des Formes), CEE Science plan (SC1 0329C), Human Frontier Science Program (RG 55-94). LG Nowak was supported by grants from the Ministère de l'Éducation Supérieure et de la Recherche and by the Fyssen Foundation.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Gross CG: **How inferior temporal cortex became a visual area.** *Cereb Cortex* 1994, 5:455-469.

An instructive and lively historical perspective on the beginnings of cortical single-unit recordings and the importance of inferotemporal cortex in vision.

2. Stone J, Dreher B, Leventhal A: **Hierarchical and parallel mechanisms in the organization of the visual cortex.** *Brain Res Rev* 1979, 1:345-394.
  3. Bullier J, Girard P, Salin PA: **The role of area 17 in the transfer of information to extrastriate visual cortex.** In *Primary Visual Cortex in Primates. Cerebral Cortex*, vol. 10. Edited by Peters A, Rockland KS. New York: Plenum; 1994:301-330.
  4. Rockland KS, Pandya DN: **Laminar origin and terminations of cortical connections of the occipital lobe in the rhesus monkey.** *Brain Res* 1979, 179:3-20.
  5. Felleman DJ, Van Essen DC: **Distributed hierarchical processing in the primate cerebral cortex.** *Cereb Cortex* 1991, 1:1-47.
  6. Girard P, Bullier J: **Visual activity in area V2 during reversible inactivation of area 17 in the macaque monkey.** *J Neurophysiol* 1989, 62:1287-1302.
  7. Schiller PH, Malpeli JG: **The effect of striate cortex cooling on area 18 cells in the monkey.** *Brain Res* 1977, 126:366-369.
  8. Girard P, Salin PA, Bullier J: **Visual activity in areas V3A and V3 during reversible inactivation of area V1 in the macaque monkey.** *J Neurophysiol* 1991, 66:1493-1503.
  9. Girard P, Salin PA, Bullier J: **Visual activity in macaque area V4 depends on area 17 input.** *Neuroreport* 1991, 2:81-84.
  10. Rocha-Miranda CE, Bender DB, Gross CG, Mishkin M: **Visual activation of neurons in inferotemporal cortex depends on striate cortex and forebrain commissures.** *J Neurophysiol* 1975, 38:475-491.
  11. Merigan WH, Nealey TA, Maunsell JHR: **Visual effects of lesions of cortical area V2 in macaques.** *J Neurosci* 1993, 13:3180-3191.
  12. Mohler CW, Wurtz RH: **Role of striate cortex and superior colliculus in visual guidance of saccadic eye movements in monkeys.** *J Neurophysiol* 1977, 40:74-94.
  13. Cowey A, Stoerig P: **Blindsight in monkeys.** *Nature* 1994, 373:247-249.
- Monkeys with lesions in V1 orient toward a visual stimulus in their blind visual field while giving a similar response as in a no-stimulus condition. This resembles the behavior of blindsight patients, who orient toward a target of which they are not conscious.
14. Girard P, Salin PA, Bullier J: **Response selectivity of neurons in area MT of the macaque monkey during reversible inactivation of area V1.** *J Neurophysiol* 1992, 67:1-10.
  15. Rodman HR, Gross CG, Albright TD: **Afferent basis of visual response properties in area MT of the macaque: I. Effects of striate cortex removal.** *J Neurosci* 1989, 9:2033-2050.
  16. Barbur JL, Watson JDG, Frackowiak RSJ, Zeki S: **Conscious visual perception without V1.** *Brain* 1993, 116:1293-1302.

17. Perenin MT: **Discrimination of motion direction in perimetricaly blind fields.** *Neuroreport* 1991, 2:397-400.
  18. Cowey A, Stoerig P: **The neurobiology of blindsight.** *Trends Neurosci* 1991, 14:140-145.
  19. Michalski A, Wimbome BM, Henry GH: **The effect of reversible cooling of cat's primary visual cortex on the responses of area 21a neurons.** *J Physiol (Lond)* 1993, 466:133-156.
  20. Guedes R, Watanabe S, Creutzfeldt OD: **Functional role of association fibres from a visual association area: the posterior suprasylvian sulcus of the cat.** *Exp Brain Res* 1983, 49:13-27.
  21. Guido W, Tong L, Spear PD: **Afferent bases of spatial- and temporal-frequency processing by neurons in the cat's postero-medial lateral suprasylvian cortex: effects of removing areas 17, 18 and 19.** *J Neurophysiol* 1990, 5:1636-1651.
  22. Sherk H: **Area 18 cell responses in cat during reversible inactivation of area 17.** *J Neurophysiol* 1978, 41:204-215.
  23. Casanova C, Michaud Y, Morin C, McKinley PA, Molotchnikoff S: **Visual responsiveness and direction selectivity of cells in area 18 during local reversible inactivation of area 17 in cats.** *Vis Neurosci* 1992, 9:581-593.
  24. Sprague JM, Levy J, DiBernardino A, Berlucchi G: **Visual cortical areas mediating form discrimination in the cat.** *J Comp Neurol* 1977, 172:441-488.
  25. Vandebussche E, Sprague JM, De Weerd P, Orban GA: **Orientation discrimination in the cat: its cortical locus. I. Areas 17, and 18.** *J Comp Neurol* 1991, 305:632-658.
  26. Raiguel SE, Lagae L, Gulyas B, Orban GA: **Response latencies of visual cells in macaque areas V1, V2 and V5.** *Brain Res* 1989, 493:155-159.
  27. Celebrini S, Thorpe S, Trotter Y, Imbert M: **Dynamics of orientation coding in area V1 of the awake primate.** *Vis Neurosci* 1993, 10:811-825.
  28. Maunsell JHR, Gibson JR: **Visual response latencies in striate cortex of the macaque monkey.** *J Neurophysiol* 1992, 4:1332-1334.
  29. Vogels R, Orban GA: **Activity of inferior temporal neurons during orientation discrimination with successively presented gratings.** *J Neurophysiol* 1994, 71:1428-1451.
  30. Nowak LG, Munk MHJ, Girard P, Bullier J: **Visual latencies in areas V1 and V2 of the macaque monkey.** *Vis Neurosci* 1995, 12:371-385.
- Latencies to visual stimulation have overlapping distributions in areas V1 and V2. V2 neurons have latencies that are on average 10 ms longer than those of V1 neurons. In V1, neurons driven by the magnocellular LGN neurons have latencies about 20 ms shorter than neurons driven by the parvocellular afferents.
31. Munk MHJ, Nowak LG, Girard P, Chounlamountri N, Bullier J: **Visual latencies in cytochrome oxidase bands of macaque area V2.** *Proc Natl Acad Sci USA* 1995, 92:988-992.
- Latencies to visual stimulation in thin cytochrome oxidase bands are longer by about 20ms than latencies in pale and thin bands. The thin cytochrome oxidase bands thus constitute a slow channel with layer 4Cβ in V1.
32. Sandell JH, Schiller PH: **Effect of cooling area 18 on striate cortex cells in the squirrel monkey.** *J Neurophysiol* 1982, 48:38-48.
  33. Maunsell JHR: **Physiological evidence for two visual subsystems.** In *Matters of Intelligence*. Edited by Vaina LM. Dordrecht: Reidel; 1987:59-87.
  34. Dinse HR, Krüger K: **The timing of processing along the visual pathway in the cat.** *Neuroreport* 1994, 5:893-897.
- This study shows that the order of activation of cortical areas in the cat does not correspond to that predicted by the hierarchical ranking of cortical areas based on the anatomy of cortico-cortical connections.
35. Bolz J, Rosner G, Wässle H: **Response latency of brisk-sustained (X) and brisk-transient (Y) cells in the cat retina.** *J Physiol (Lond)* 1982, 328:171-190.
  36. Shapley R: **Visual sensitivity and parallel retinocortical channels.** *Annu Rev Psychol* 1990, 41:635-658.
  37. Merigan WH, Maunsell JHR: **How parallel are the primate visual pathways?** *Annu Rev Neurosci* 1993, 16:369-402.

38. Casagrande VA: **A third visual pathway to primate area V1.** *Trends Neurosci* 1994, 17:305–310.
39. Hendry SHC, Yoshioka TY: **A neurochemically distinct third channel in the macaque dorsal lateral geniculate nucleus.** *Science* 1994, 264:575–577.
40. Schiller PH, Logothetis NK, Charles ER: **Role of the color-opponent and broad-band channels in vision.** *Vis Neurosci* 1990, 5:321–346.
41. Hubel DH, Livingstone MS: **Segregation of form, color, and stereopsis in primate area 18.** *J Neurosci* 1987, 7:3378–3415.
42. Lachica EA, Beck PD, Casagrande VA: **Parallel pathways in macaque monkey striate cortex: anatomically defined columns in layer III.** *Proc Natl Acad Sci USA* 1992, 89:3566–3570.
43. Lachica EA, Beck PD, Casagrande VA: **Intrinsic connections of layer III of striate cortex in squirrel monkey and bushy monkey: correlations with patterns of cytochrome oxidase.** *J Comp Neurol* 1993, 329:163–187.
44. Yoshioka T, Levitt JB, Lund JS: **Independence and merger of thalamocortical channels within macaque monkey primary visual cortex: anatomy of interlaminar projections.** *Vis Neurosci* 1994, 11:467–489.
- Demonstrates the convergence of M and P signals in interblob regions of layers 2 and 3 of V1. Shows the convergence of M and P pathways in the cytochrome oxidase blobs and that P signals have to relay in layer 4A to reach the blobs.
45. Levitt JB, Yoshioka T, Lund JS: **Intrinsic cortical connections in macaque visual area V2: evidence for interaction between different functional streams.** *J Comp Neurol* 1994, 342:551–570.
- Shows a dense array of interconnections between the different cytochrome oxidase bands of V2. This suggests that there are more interactions between the channels than assumed in the parallel pathways model.
46. Malach R, Tootell RGB, Malonek D: **Relationship between orientation domains, cytochrome oxidase stripes, and intrinsic horizontal connections in squirrel monkey area V2.** *Cereb Cortex* 1994, 4:151–165.
- This study used a combination of optical imaging of cortical activity, tract tracing and cytochrome oxidase histochemistry to reveal the specificity and interconnections of functional domains. It shows a large array of interconnections between the cytochrome oxidase stripes in V2. It also demonstrates that interconnections within V2 are not restricted to the domains with the same optimal orientations.
47. Malpeli JG, Schiller PH, Colby CL: **Response properties of single cells in monkey striate cortex during reversible inactivation of individual geniculate laminae.** *J Neurophysiol* 1981, 46:1102–1119.
48. Nealey TA, Maunsell JHR: **Magnocellular and parvocellular contributions to the responses of neurons in macaque striate cortex.** *J Neurosci* 1994, 14:2069–2079.
- Physiological demonstration that neurons in all cytochrome oxidase subunits in V1 receive convergent inputs from magnocellular and parvocellular LGN layers.
49. DeYoe EA, Van Essen DC: **Segregation of efferent connections and receptive field properties in visual area V2 of the macaque.** *Nature* 1985, 317:58–61.
50. Levitt JB, Kiper DC, Movshon JA: **Receptive fields and functional architecture of macaque V2.** *J Neurophysiol* 1994, 71:2517–2542.
- Quantitative study of the receptive field properties of neurons in different cytochrome oxidase bands in V2. Demonstrates that color, length and orientation selectivities are less segregated in different cytochrome oxidase bands than previously thought.
51. Ferrera VP, Nealey TA, Maunsell JHR: **Response in macaque visual area V4 following inactivation of the parvocellular and magnocellular LGN pathways.** *J Neurosci* 1994, 14:2080–2088.
- Demonstrates directly that V4 neurons are driven by inputs from magnocellular and parvocellular neurons of the LGN relayed in V1 and V2. This goes against the notion that V4 processes exclusively parvocellular-type information.
52. Cheng K, Hasegawa T, Saleem KS, Tanaka K: **Comparison of neuronal selectivity for stimulus speed, length, and contrast in the prestriate visual cortical areas V4 and MT of the macaque monkey.** *J Neurophysiol* 1994, 71:2269–2280.
- Shows more similarities between the response selectivities of neurons in areas V4 and MT than assumed by the model of functional specialization of cortical areas.
53. Maunsell JHR, Nealey TA, Depriest DD: **Magnocellular and parvocellular contributions to responses in the middle temporal visual area (MT) of the macaque monkey.** *J Neurosci* 1991, 10:3323–3334.
54. Gegenfurtner KR, Kiper DC, Beusmans JMH, Carandini M, Zaidi Q, Movshon JA: **Chromatic properties of neurons in macaque MT.** *Vis Neurosci* 1994, 11:455–466.
55. Saito H, Tanaka K, Isono H, Yasuda M, Mikami A: **Directionally selective response of cells in the middle temporal area (MT) of the macaque monkey to the movement of equiluminous opponent color stimuli.** *Exp Brain Res* 1989, 75:1–14.
56. Livingstone MS, Hubel DH: **Anatomy and physiology of a color system in the primate visual cortex.** *J Neurosci* 1984, 4:309–356.
57. T'so DY, Gilbert CD: **The organization of chromatic and spatial interactions in the primate striate cortex.** *J Neurosci* 1988, 8:1712–1727.
58. Lennie P, Krauskopf J, Sclar G: **Chromatic mechanisms in striate cortex of macaque.** *J Neurosci* 1990, 10:649–669.
59. Peterhans E, Von der Heydt R: **Functional organization of area V2 in the alert macaque.** *Eur J Neurosci* 1993, 5:509–524.
60. Schiller PH: **The effects of V4 and middle temporal (MT) lesions on visual performances in the rhesus monkey.** *Vis Neurosci* 1993, 10:717–746.
61. Pasternak T, Merigan WH: **Motion perception following lesions of the superior temporal sulcus in the monkey.** *Cereb Cortex* 1994, 4:247–259.
- Large lesions of the superior temporal sulcus (restricted to the gray matter) produce limited effects on motion-related tasks. The visual cortex can process motion without the middle temporal area and its surrounding areas.
62. Heywood CA, Gadotti A, Cowey A: **Cortical area V4 and its role in the perception of color.** *J Neurosci* 1992, 12:4056–4065.
63. Newsome WT, Paré EB: **A selective impairment of motion processing following lesions of the middle temporal visual area (MT).** *J Neurosci* 1988, 8:2201–2211.
64. Lomber SG, Cornwell P, Sun J-S, MacNeil MA, Payne BR: **Reversible inactivation of visual processing operations in middle suprasylvian cortex of the behaving cat.** *Proc Natl Acad Sci USA* 1994, 91:2999–3003.
- Shows a strong, specific and reversible effect of inactivation of the lateral suprasylvian sulcus on visual discrimination of targets masked by dynamic visual noise. This technique by-passes the difficulties of interpretation related to the extensive post-operative recovery of lesion studies.
65. Ullman S: **Sequence seeking and counter streams: a computational model for bidirectional information flow in the visual cortex.** *Cereb Cortex* 1995, 5:782–792.
66. Mumford D: **Neuronal architectures for pattern-theoretic problems.** In *Large Scale Neuronal Theories of the Brain*. Edited by Koch C, Davis J. Cambridge, Massachusetts: MIT Press; 1994:125–152.
67. Tononi G, Sporns O, Edelman GM: **Reentry and the problem of integrating multiple cortical areas: simulation of dynamic integration in the visual system.** *Cereb Cortex* 1992, 2:310–335.
68. Dupont P, Orban GA, De Bruyn B, Verbruggen A, Mortelmans L: **Many areas in the human brain respond to visual motion.** *J Neurophysiol* 1994, 72:1420–1424.
69. Gulyas B, Roland PE: **Processing and analysis of form, colour and binocular disparity in the human brain: functional anatomy by positron emission tomography.** *Eur J Neurosci* 1994, 6:1811–1828.
70. Salin P-A, Bullier J: **Corticocortical connections in the visual system: structure and function.** *Physiol Rev* 1995, 75:107–154.

J Bullier, Cerveau et Vision, INSERM 371, 18 avenue du doyen Lépine, 69500 Bron/Lyon, France.

J Bullier E-mail: bullier@cismibm.univ-lyon1.fr

LG Nowak, Section of Neurobiology, Yale University School of Medicine, New Haven, Connecticut 06510, USA.